

Parkinson disease among patients treated for benign prostatic hyperplasia with α 1 adrenergic receptor antagonists

Rahul Sasane, ... , John J. Renger, David J. Stone

J Clin Invest. 2021. <https://doi.org/10.1172/JCI145112>.

Clinical Medicine In-Press Preview Genetics Neuroscience

BACKGROUND. Recently the α 1 adrenergic receptor antagonist terazosin was shown to activate PGK1, a possible target for the mitochondrial deficits in Parkinson disease related to its function as the initial enzyme in ATP synthesis during glycolysis. An epidemiologic study of terazosin users showed a lower incidence of Parkinson disease when compared to users of tamsulosin, an α 1 adrenergic receptor antagonist of a different class that does not activate PGK1. However, prior research on tamsulosin has suggested that it may in fact potentiate neurodegeneration, raising the question of whether it is an appropriate control group.

METHODS. To address this question, we undertook an epidemiological study on Parkinson disease occurrence rate in 113,450 individuals from the U.S.A. with > 5 years of follow-up. Patients were classified as tamsulosin users (n = 45,380), terazosin/alfuzosin/doxazosin users (n = 22,690) or controls matched on age, gender and Charlson Comorbidity Index score (n = 45,380).

RESULTS. Incidence of Parkinson disease in tamsulosin users was 1.53%, which was significantly higher than that in both terazosin/alfuzosin/doxazosin users (1.10%; $p < 0.0001$) and matched controls (1.01%; $p < 0.0001$). Terazosin/alfuzosin/doxazosin users did not differ in Parkinson disease risk from [...]

Find the latest version:

<https://jci.me/145112/pdf>



Parkinson Disease among Patients Treated for Benign Prostatic Hyperplasia with α_1 Adrenergic Receptor Antagonists

Rahul Sasane,¹ Amy Bartels,² Michelle Field,² Maria I Sierra,² Sridhar Duvvuri,¹ David Gray,¹ Sokhom Pin,¹ John J Renger,¹ and David J Stone^{1,3}

1. Cerevel Therapeutics, Cambridge, MA, USA

2. Optum Life Sciences, Eden Prairie, MN, USA

3. Corresponding Author: David J Stone, Cerevel Therapeutics, LLC, 222 Jacobs Street. Suite 200, Cambridge, MA 02141. Phone +1.857.299.7114; email david.stone@cerevel.com

Conflict of Interest

Disclosures: Rahul Sasane, Sridhar Duvvuri, David Gray, Sokhom Pin, John J Renger, and David J Stone were employees of and equity shareholders in Cerevel Therapeutics LLC at the time this work was conducted. At the time of the study, Amy Bartels, Michelle Field, and Maria I Sierra were employed by Optum, which was contracted by Cerevel to perform the analyses. Employment of none of the authors was contingent upon the publication of the manuscript. The authors declare no other conflicts of interest.

Abstract

Background. Recently the α_1 adrenergic receptor antagonist terazosin was shown to activate PGK1, a possible target for the mitochondrial deficits in Parkinson disease related to its function as the initial enzyme in ATP synthesis during glycolysis. An epidemiologic study of terazosin users showed a lower incidence of Parkinson disease when compared to users of tamsulosin, an α_1 adrenergic receptor antagonist of a different class that does not activate PGK1. However, prior research on tamsulosin has suggested that it may in fact potentiate neurodegeneration, raising the question of whether it is an appropriate control group.

Methods. To address this question, we undertook an epidemiological study on Parkinson disease occurrence rate in 113,450 individuals from the U.S.A. with ≥ 5 years of follow-up. Patients were classified as tamsulosin users (n = 45,380), terazosin/alfuzosin/doxazosin users (n = 22,690) or controls matched on age, gender and Charlson Comorbidity Index score (n = 45,380).

Results. Incidence of Parkinson disease in tamsulosin users was 1.53%, which was significantly higher than that in both terazosin/alfuzosin/doxazosin users (1.10%; $p < 0.0001$) and matched controls (1.01%; $p < 0.0001$).

Terazosin/alfuzosin/doxazosin users did not differ in Parkinson disease risk from matched controls ($p = 0.29$).

Conclusions. These results suggest that zosins may not confer a protective effect against Parkinson disease, but rather that tamsulosin may in some way potentiate Parkinson disease progression.

Funding: This work was supported by Cerevel Therapeutics.

INTRODUCTION

Parkinson disease (PD) is the second most common neurodegenerative disorder after Alzheimer disease. Due to the aging of the world population, the incidence of PD is increasing with 6.1 million affected individuals in 2016, as compared to 2.5 million affected individuals in 1990 (1). Development of disease-modifying therapies is the greatest unmet need in treatment of neurodegenerative disease; however, the high cost and long timeline of drug development limit the number of therapeutic hypothesis that can be rigorously evaluated in clinical studies (2). Furthermore, when an ineffective approach is pursued, this is often not apparent until completion of large phase 2 or 3 trials. Therefore, early identification of treatment targets with increased probability of success is critical to minimize negative resource and capital opportunity cost of failure, unproductive allocation of scarce resources of time and capital, and to increase the speed to finding effective therapeutics for patients with significant unmet need.

Recent evidence from animal models and retrospective epidemiological studies has suggested that phosphoglycerate kinase 1 (PGK1) is a target of interest for disease modification in PD (3). PGK1 is the first enzyme in ATP generation via glycolysis, and therefore PGK1 potentiation may ameliorate cellular pathophysiology in disorders with an established mitochondrial deficit/phenotype such as PD through increasing ATP production in the context of reduced cellular energy production. In 2015, terazosin, an α -1 adrenergic receptor antagonist frequently prescribed for benign prostatic hyperplasia (BPH), was found to bind to PGK1 and potentiate its kinase activity (4). This PKG1 pharmacology has also been demonstrated in related drugs harboring quinazoline motifs, such as alfuzosin and doxazosin (4). Furthermore, an analysis of the Truven database and Danish nationwide health registries demonstrated that individuals treated with terazosin, alfuzosin, or doxazosin showed lower rates of both the development of PD and PD-related diagnoses when compared to patients treated with another BPH treatment, tamsulosin, that does not interact with PGK1 (3,5). To that extent, a clinical study evaluating the safety and tolerability of terazosin, 5 mg once daily for 12 weeks, in patients with Parkinson's disease (H&Y stage I-III) has been initiated ([NCT03905811](#)).

While the prospect that terazosin may have a protective effect in PD is promising, it is important to closely review the evidence for possible alternative explanations for the observed risk imbalance. Notably, in the analysis by Cai et al, there was no control group of age-matched patients who were not on a BPH treatment. Therefore, while a protective

Sasane et al. PD among alpha 1 Adrenergic Receptor Agonists w BPH effect associated with terazosin use is possible, an alternative interpretation could be an increased incidence or rate of progression of PD in patients using tamsulosin as the comparison group (3). This idea is not without precedent, Duan et and colleagues found a positive correlation between tamsulosin use and dementia risk when compared to terazosin, doxazosin, alfuzosin, or no BPH medication (6). The inclusion of a propensity-score matched (PSM) no-BPH drug cohort is critical for interpretation of differences between drug effects (i.e. protection via terazosin vs. increased risk via tamsulosin

To further study the implications of tamsulosin and terazosin use on PD risk, we designed a large, retrospective study looking at the incidence of PD in 3 non-overlapping groups of patients: those treated with terazosin, alfuzosin, or doxazosin (Zosin Cohort); those treated with tamsulosin (Tamsulosin Cohort); and an appropriately matched Control Cohort with no BPH drug treatment.

RESULTS

This epidemiological study on risk of PD included 113,450 individuals with ≥ 5 years of follow-up. Figure 1 shows the sample selection process for the Zosin and Tamsulosin Cohorts, and the Control Cohort, based upon defined inclusion and exclusion criteria. Propensity score matching (PSM) was used to balance the cohorts on demographic and clinical differences (7). Before the PSM procedure was conducted, the initial sample of patients taking tamsulosin was 87,674 and 35,405 were taking a zosin medication (7). From among the 2,151,212 patients identified as eligible controls, the SAS survey select procedure was used to select a random sample of 811,587 patients, stratified by age groups corresponding to the Zosin Cohort after matching to the Tamsulosin Cohort (8).

Propensity Score Matching Results

As a result of the PSM procedure, the final Zosin Cohort included 22,690 patients, and the Tamsulosin and Control Cohorts each included 45,380 patients (Table 1 and Table 2). Before matching, the Zosin Cohort was older, had fewer males, and lower Charlson Comorbidity Index (CCI) scores than the Tamsulosin Cohort (9). Similarly, pre-matching, the Zosin Cohort was older, had more males, and higher CCI scores than the Control Cohort.

PD Diagnoses during Follow-Up Period

As shown in Table 3, diagnoses of PD were observed more frequently in the Tamsulosin Cohort than the Zosin Cohort (1.52% vs. 1.10%, respectively, all patients; $p < 0.0001$), but they were also observed more frequently in the Tamsulosin Cohort as compared with the Control Cohort (1.53% vs. 1.01%, respectively, all patients, $p < 0.001$). The rate observed here in the Control Cohort is consistent with that seen in other published studies (10).

The Zosin Cohort did not differ in PD risk from matched controls when examining the entire cohort ($p = 0.29$) or when the analysis was restricted to those over 60 years of age ($p = 0.60$). In contrast, PD diagnoses were significantly more frequent among the Tamsulosin Cohort (1.53%) as compared with the matched Control Cohort (1.01%; $p < 0.0001$) and the Zosin Cohort (1.10%; $p < 0.0001$), and this result held when analysis was restricted to those over the age of 60 (1.78% for tamsulosin vs. 1.24% for zosins, $p < 0.0001$; and vs. 1.19% for matched controls, $p < 0.0001$). There was some evidence of an increased frequency of PD diagnoses in those under the age of 60 for individuals taking either tamsulosin

Sasane et al. PD among alpha 1 Adrenergic Receptor Agonists w BPH (p = 0.0164) or a zosin (p = 0.0115) when compared to matched controls; however this result should be taken with caution due to the small number of patients diagnoses with PD in this age group (n = 9 controls, 15 zosin, 23 tamsulosin).

DISCUSSION

Cai et al., using the Truven database, investigated whether patients taking terazosin, alfuzosin, or doxazosin had similar proportions of patients who developed PD as compared with a tamsulosin control group (3). Their finding that patients taking a zosin medication had fewer PD diagnoses than those taking tamsulosin was significant, however their follow-up time was less than 1 year. By extending follow-up to 5 years, the current study enables examination of PD diagnoses in the longer term, which increases confidence in the diagnosis. Approximately 20% of diagnoses of PD are incorrect within the first year, but this proportion decreases to near zero after a few years. Also, their study did not have a BPH medication-free control group. Because our study included comparisons to matched control subjects, our results suggest that zosins may not confer a protective effect against PD, as proposed by Cai et al., but rather that tamsulosin may in some way potentiate neurodegeneration, in particular among patients over age 61.

This suggestion is supported by a database study by Duan et al. (6) which found the risk of dementia (not specifically in association with PD) was higher among patients taking tamsulosin, as compared with a control cohort which took no BPH medication as well as individual cohorts taking alternative BPH medications, including doxazosin, terazosin, and alfuzosin, specifically among patients over 65 years of age. The finding was significant when analyses controlled for comorbidity burden and use of chronic medications known to increase the risk of developing dementia. Thus, use of tamsulosin appears to increase the risk of cognitive adverse effects, including the development of dementia, which may not have been observed in a relatively short follow-up in clinical trials of tamsulosin or the study by Cai et al (3).

Limitations of the current study include matching only on age, gender, and CCI score. Although CCI was considered a proxy for health conditions, it would be useful to match on specific conditions, lifestyle choices (such as smoking), socioeconomic factors, and medications that are likely to affect the risk of developing PD. Furthermore, because the drugs in this study are primarily prescribed for BPH, the cohort was almost entirely male and finding therefore can only be applied to men. Retrospective observational studies have limitations and claims data do not include all possible outcomes variables permitting more in-depth analyses. The current study determined the risk of developing PD based solely on diagnosis codes; coding may be subject to error and reimbursement bias as claims are not created for the purposes of research, but for administration and billing. The results obtained are specific to patients with the type of

Sasane et al. PD among alpha 1 Adrenergic Receptor Agonists w BPH insurance included; they may not be applicable to different populations. Importantly, this study points out that because of these limitations in retrospective studies, careful attention to inclusion of all necessary control groups is essential to correctly understand the implications of any results. Because findings such as these can influence the decisions made by patients and their physicians concerning their care, it is critical that alternative interpretations are fully assessed.

This study challenges the notion that PGK1 may be a validated target for disease modification in PD and raises the question as to the best choice for BPH medications for long-term use by elderly patients. A mechanism explaining the differences we and others have observed on the association of tamsulosin use and neurodegeneration may be further elucidated by longitudinal real-world evidence, controlling for more variables associated with risk of PD and dementia.

METHODS

Data Source

The Optum Research Database (ORD), Optum's proprietary administrative claims research database composed of pharmacy and medical claims. The ORD is a proprietary administrative claims research database, with de-identified and HIPAA compliant medical and pharmacy claims data from 1993-present on more than 73 million lives. In 2018, approximately 19% of the US commercially enrolled population, 21% of the Medicare Advantage population and 22% of the Medicare Part D (with pharmacy claims only) populations were represented in the ORD.

Study Design

This was a retrospective observational study including claims data for the period January 1, 2010 to December 31, 2019 (study period). An index event was defined by the first prescription fill for an alpha blocker medication of interest for BPH (terazosin, doxazosin, afluzosin, tamsulosin) during the identification period (see Supplemental Figure S1) January 1, 2011 to December 31, 2019. A baseline period of at least 12 months with continuous medical and pharmacy benefits prior to the index event was set to identify any previous PD diagnosis or PD medication (DOPA decarboxylase/DA precursor, dopamine agonists, MAO B inhibitors, COMT inhibitors, anticholinergics, adamantanes, adenosine A2a agonists; see Supplemental Table S1). A follow-up period of at least 5 years of continuous medical and pharmacy benefits post-index was used to assess PD diagnosis occurring after the initiation of treatment with the index medication.

Study Sample

The study included commercially and Medicare Advantage Part-D (MAPD) insured patients (≥ 18 years of age) taking medications of interest for BPH (terazosin, doxazosin, afluzosin, tamsulosin), who had no evidence of PD during the baseline period. Three mutually exclusive cohorts were created based upon presence or absence of BPH medication of interest, along with other criteria: the Zosin Cohort taking terazosin, doxazosin, or alfuzosin; the Tamsulosin Cohort, which was only taking tamsulosin; and the Control Cohort, which was taking none of the 4 drugs of interest. For all

Sasane et al. PD among alpha 1 Adrenergic Receptor Agonists w BPH cohorts, continuous enrollment with medical and pharmacy benefits for the entire study period, including 5 years of follow-up, were required.

Patients were further included in the Zosin or Tamsulosin Cohorts by at least 2 prescription fills in the same year for a zosin (terazosin, doxazosin, afluzosin) or tamsulosin. They were excluded by the presence of claims containing PD diagnoses codes (ICD code 332 or G20) or 2 or more PD medication fills in the baseline period.

Patients were included in the Control Cohort with at least 1 prescription fill for any prescription, other than the 4 BPH medications of interest during the study period. A random index date was chosen as a date of a prescription. They were excluded by PD diagnosis codes (ICD code 332 or G20) or 2 or more PD medication fills in the baseline period, or fills for terazosin, doxazosin, alfuzosin, or tamsulosin.

Measures/Outcomes

Patient characteristics were noted, including age as of the index date, gender, and Charlson Comorbidity Index (CCI) score at index (9). Mean (\pm standard deviation [SD]) continuous age; n (%) male gender; and mean (SD) continuous CCI score were compared. During the follow-up period, the number (%) of patients with PD diagnosis was identified by ICD-9/10 codes 332 or G20 and compared between Zosin and Tamsulosin Cohorts, and between Zosin and Control Cohorts, stratified by age group (all, 18-60 years, and 61+ years).

Analytical Methods

Randomization of Controls

From among the 2,151,212 patients identified as eligible controls, the SAS survey select procedure (8) was used for a simple random sample. We stratified by the age group at index (18 to 50, 51 to 55, 56 to 60, 61 to 65, 66 to 70, 71 to 75, 76 to 80, 81+). For the sampling rate, we used the age group distribution of the Zosin Cohorts after matching to the Tamsulosin Cohort. The final random sample of controls included 811,587 patients.

PSM Matching Procedure

Propensity score matching (PSM) was used to balance the cohorts on demographic and clinical differences including age on index date, gender, and mean CCI score (7). A greedy match procedure provided 2 control matches per zosin case. First, patients taking zosins were matched to patients taking tamsulosin. From that matching output, the zosin-matched patients were matched to randomized controls using the same criteria. For any cases without 2 controls, cases and all corresponding controls were dropped.

Statistics

Statistical analyses were conducted comparing matched Zosin Cohort vs Control Cohort, Zosin vs. Tamsulosin Cohort, and Tamsulosin vs. Control Cohort, with the SAS procedure, proc genmod, comparing patient characteristics used for the PSM procedure (7,8). Cohorts were compared using Chi-square statistics for outcomes of PD diagnoses over the 5-year minimum follow-up period; $p < 0.05$ was considered significant.

Study Approval

Only completely de-identified data were accessed for this study and its conduct aligned with US Health Insurance Portability and Accountability Act regulations. Thus, Institutional Review Board approval was neither obtained nor sought.

Author contributions

DJS: Initial conception of study, analysis and MS preparation; RS: Initial conception of study, analysis and MS preparation; DG: Contribution to initial concept for study and to MS preparation (review/revision); SD: Contribution to initial concept for study and to MS preparation (review/revision); SP: Contribution to initial concept for study and to MS preparation (review/revision); JJR: Contribution to initial concept for study and to MS preparation (review/revision); AB:

Contribution to study design, data collection and analysis, and MS preparation; MF: Contribution to study design, data analysis, and MS preparation; MIS: Contribution to study design, data analysis, and MS preparation.

Acknowledgments


Medical writing support was provided by Caroline Jennermann, of Optum HEOR, as funded by Cerevel. Armen Zakharyan was responsible for development of the models used in this work. We thank Monica Frazer for contributions to the development of the manuscript; Phil Iredale, Yong Mi Choi, Georgette Suidan, David Zhang, Kimberly Kerr, and Hanh Nguyen for helpful discussion and for contributions to the design of the study; and Timothy Bancroft for analytical guidance.

REFERENCES

1. Global Burden of Disease Causes of Death Collaborators. Global, regional, and national ag-sex specific mortality for 264 causes of death, 1980-2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 2017;390(10100):1151-1210.
2. Herper M. How much does pharmaceutical innovation cost? A look at 100 companies. *Forbes Healthcare* 2013, Aug 11. Forbes Media LLC
3. Cai R, Zhang Y, Simmering JE, et al. Enhancing glycolysis attenuates Parkinson's disease progression in models and clinical databases. *J Clin Invest* 2019;129(10):4539-4549.
4. Chen X, Zhao C, Lie X, et al. Terazosin activates pgk1 and Hsp90 to promote stress resistance. *Nat Chem Biol* 2015;11(1):19-25.
5. Simmering JE, Welsh MJ, Liu L, Narayanan NS, Pottegård A. Association of glycolysis-enhancing α -1 blockers with risk of developing Parkinson disease. *JAMA Neurol*. Doi:10.1001/jamaneurol.2020.5157
6. Duan Y, Grady JJ, Albertsen PC, Wu ZH. Tamsulosin and the risk of dementia in older men with benign prostatic hyperplasia. *Pharmacoepidemiol Drug Saf*. 2018;27(3):340-348.
7. Rosenbaum PR, Rubin DB. The central role of the propensity score in observational studies for causal effects. *Biometrika* 1983. 70(1):41-55.
8. SAS Institute, Inc. 2020. SAS Enterprise Guide software version 7.15
9. Quan H, Li B, Couris CM, et al. Updating and validating the Charlson comorbidity index and score for risk adjustment in hospital discharge abstracts using data from 6 countries. *Am J Epidemiol* 2011;173(6):676-682.
10. Hirsch L, Jette N, Frolkis A, Steeves T, Pringsheim T. The incidence of Parkinson's disease: a systematic review and meta-analysis. *Neuroepidemiology* 2016;46(4):292-300.

Figure 1. Initial Sample Selection and Attrition

A. Zosin and Tamsulosin Cohorts



B. Control Cohort




Table 1. Age, sex, and comorbidity score distribution in Zosin to Tamsulosin Cohorts before and after propensity score matching

	Pre-Matching				Post-Matching			
	Zosin (N=35,405)	Tamsulosin (N= 87,674)	Std Diff %	P	Zosin (N=22,690)	Tamsulosin (N= 45,380)	Std Diff %	P
Age Mean (SD)	71.48 (9.53)	69.78(10.55)	16.97	<.0001	70.33 (9.92)	70.33 (9.92)	0.00	1.000
Age Group, Years % (N)								
18 to 60	13.87 (4,912)	18.84 (16,515)			17.08 (3,876)	17.08 (7,752)		
61 to 65	6.06 (2,147)	6.54 (5,734)			6.74 (1,529)	6.74 (3,058)		
66 to 70	18.97 (6,715)	20.17 (17,681)			19.94 (4,524)	19.94 (9,048)		
71 to 75	21.01 (7,438)	21.28 (18,654)			20.45 (4,641)	20.45 (9,282)		
76 to 80	18.74 (6,636)	16.00 (14,031)			18.22 (4,134)	18.22 (8,268)		
81+	21.34(7,557)	17.18 (15,059)			17.57 (3,986)	17.57 (7,972)		
Male, % (n)	78.37 (27,747)	96.38 (84,499)	-56.33	<.0001	98.83 (22,424)	98.83 (44,848)	0.00	1.000
CCI score, mean (SD)	1.095 (1.491)	1.139 (1.573)	-2.85	<.0001	1.000 (1.399)	1.000 (1.399)	0.00	1.000
CCI score categories, % (n)								
0-1	73.83 (26,138)	71.76 (62,915)			71.11 (16,134)	71.11 (32,268)		
2 to 3	19.62 (6,945)	20.81 (18,245)			22.41 (5,084)	22.41 (10,168)		
4 to 5	5.04 (1,784)	5.44 (4,772)			5.12 (1,161)	5.12 (2,322)		
≥6	1.52 (538)	1.99 (1,742)			1.37 (311)	1.37 (622)		

Table 2. Characteristics of patients comparing Zosin to Control Cohorts before and after PSM

	Pre-Matching				Post-Matching			
	Zosin (N=35,405)	Control (N=811,587)	Std Diff %	P	Zosin (N=22,690)	Control (N=45,380)	Std Diff %	P
Age Mean (SD)	71.48 (9.53)	70.43 (11.23)	10.08	<.0001	70.33 (9.92)	70.33 (9.92)	0.00	1.000
Age Group, Years % (N)								
18 to 60	13.87 (4,912)	16.53 (134,152)			17.08 (3,876)	17.08 (7,752)		
61 to 65	6.06 (2,147)	6.67 (54,167)			6.74 (1,529)	6.74 (3,058)		
66 to 70	18.97 (6,715)	19.23 (156,043)			19.94 (4,524)	19.94 (9,048)		
71 to 75	21.01 (7,438)	20.21 (164,048)			20.45 (4,641)	20.45 (9,282)		
76 to 80	18.74 (6,636)	17.03 (138,247)			18.22 (4,134)	18.22 (8,268)		
81+	21.34(7,557)	20.32 (164,930)			17.57 (3,986)	17.57 (7,972)		
% Male	78.37 (27,747)	31.17 (252,952)	107.72	<.0001	98.83 (22,424)	98.83 (44,848)	0.00	1.000
CCI score, mean (SD)	1.095 (1.491)	0.837 (1.345)	18.16	<.0001	1.000 (1.399)	1.000 (1.399)	0.00	1.000
CCI score categories, % (n)								
0-1	73.83 (26,138)	79.38 (644,266)			71.11 (16,134)	71.11 (32,268)		
2 to 3	19.62 (6,945)	16.15 (131,062)			22.41 (5,084)	22.41 (10,168)		
4 to 5	5.04 (1,784)	3.24 (26,300)			5.12 (1,161)	5.12 (2,322)		
≥6	1.52 (538)	1.23 (9,959)			1.37 (311)	1.37 (622)		

Table 3. Proportion of Cohorts with PD Diagnosis

Age Groups	Zosin Cohort (n=22,690)		Control Cohort (n=45,380)		Zosin: Control Comparison	
	Total Patients	With PD, n(%)	Total Patients	With PD, n(%)	ChiSq	p Value
All Patients	22,690	249 (1.10)	45,380	458 (1.01)	1.13	0.2878
Age 18 - 60	3,876	15 (0.39)	7,752	9 (0.12)	6.39	0.0115
Age 61+	18,814	234 (1.24)	37,628	449 (1.19)	0.27	0.6047

Age Groups	Tamsulosin Cohort (n=45,380)		Control Cohort (n=45,380)		Tamsulosin: Control Comparison	
	Total Patients	With PD, n(%)	Total Patients	With PD, n(%)	ChiSq	p Value
All Patients	45,380	693 (1.53)	45,380	458 (1.01)	48.66	<.0001
Age 18 - 60	7,752	23 (0.30)	7,752	9 (0.12)	5.76	0.0164
Age 61+	37,628	670 (1.78)	37,628	449 (1.19)	44.36	<.0001

Age Groups	Tamsulosin Cohort (n=45,380)		Zosin Cohort (n=22,690)		Tamsulosin: Zosin Comparison	
	Total Patients	With PD, n(%)	Total Patients	With PD, n(%)	ChiSq	p Value
All Patients	45,380	693 (1.53)	22,690	249 (1.10)	22.62	<.0001
Age 18 - 60	7,752	23 (0.30)	3,876	15 (0.39)	0.58	0.4477
Age 61+	37,628	670 (1.78)	18,814	234 (1.24)	25.57	<.0001

Supplemental Table S1. Medications of Interest

Medication class (USC class level)	Medication (generic)
BPH medications	
Alpha blockers	terazosin, doxazosin, afluzosin, tamsulosin
PD medications	
DOPA decarboxylase inhibitor/ DA precursor	levodopa, carbidopa, carbidopa-levodopa
Dopamine agonists	pramipexole, ropinirole, rotigotine, apomorphine, bromocriptine mesylate
MAO B inhibitors	selegiline, rasagiline mesylate, safinamide mesylate
COMT inhibitors	entacapone, tolcapone, carbidopa-levodopa-entacapone
Anticholinergics	trihexyphenidyl, benztropine
Adamantanes	amantadine
Adenosine A2a antagonists	istradefylline

Note. COMT=catechol O-methyltransferase; DA=dopamine agonist; DOPA=dihydroxyphenylalanine; MAO=monoamine oxidase; PD=Parkinson disease

Supplemental Figure S1. Study Design

